

PRESCRIBER START FORM - GENERALIZED MYASTHENIA GRAVIS (gMG)



FAX: 1.800.420.5150



MAIL: 100 College Street
New Haven, CT 06510



PHONE: 1.888.765.4747
8:30 AM to 8 PM ET Monday-Friday



EMAIL: OneSource@Alexion.com



OneSource™ is a complimentary, personalized patient support program offered by Alexion. It's designed to support patients' specific needs throughout treatment. For more information, visit www.AlexionOneSource.com. Contact OneSource if you have any questions while completing the forms.



INSTRUCTIONS FOR HEALTHCARE PROFESSIONALS:

To enroll your patient in OneSource, please follow these steps:

- 1 Have your patient complete all required sections and read the Authorization to Share Health Information on the **Patient Services Enrollment Form**
- 2 Complete all required sections on **PAGE 1**
- 3 Sign the Prescriber Certification on **PAGE 2**
- 4 **FAX PAGES 1-2 of the completed form and copies of the front and back of the patient's medical insurance and pharmacy coverage cards to OneSource.** If applicable, **fax the Vaccination Order Form (PAGE 3)** to OneSource as well.

Fields in red with asterisks are required.*

STEP 1: PATIENT INFORMATION

PATIENT NAME (FIRST, LAST)*	DATE OF BIRTH (MM/DD/YYYY)*	PATIENT PHONE NUMBER*	PATIENT EMAIL
-----------------------------	-----------------------------	-----------------------	---------------

STEP 2: CLINICAL DIAGNOSIS

SOLIRIS and ULTOMIRIS are FDA approved for antibody-positive status. If a payer requires prior authorization and/or has a clinical policy, they may require proof of antibody status.

INDICATION (check one)*: GENERALIZED MYASTHENIA GRAVIS (gMG)	ANTIBODY STATUS (check one)*:
<input type="checkbox"/> ICD-10: G70.00 Myasthenia gravis without (acute) exacerbation <input type="checkbox"/> ICD-10: G70.01 Myasthenia gravis with (acute) exacerbation	<input type="checkbox"/> ANTI-AChR ANTIBODY POSITIVE (gMG) <input type="checkbox"/> UNKNOWN (CONTACT ONESOURCE FOR QUESTIONS)

STEP 3: INSURANCE INFORMATION

Complete this section **OR** attach copies of patient's medical and pharmacy insurance card(s).*

PLEASE PROVIDE SUMMARY OF BENEFIT INVESTIGATION FOR ULTOMIRIS AND SOLIRIS

<input type="checkbox"/> COPIES OF PATIENT'S INSURANCE CARD(S) ATTACHED <input type="checkbox"/> PATIENT DOES NOT HAVE INSURANCE	PRIMARY MEDICAL INSURANCE	SECONDARY MEDICAL INSURANCE	PHARMACY COVERAGE
INSURANCE PROVIDER*			
INSURANCE PHONE #*			
CARDHOLDER NAME*			
CARDHOLDER DATE OF BIRTH*			
MEMBER ID*			
POLICY #*			
GROUP #*			
BIN/PCN #			

STEP 4: HEALTHCARE PRESCRIBER INFORMATION

FIRST NAME*	LAST NAME*	PROVIDER EMAIL*
ADDRESS*		PHONE NUMBER*
CITY*	STATE*	ZIP*
PRACTICE NAME	TAX ID #*	NPI #*
OFFICE CONTACT NAME	EMAIL	FAX NUMBER*

STEP 5: SITE OF CARE

SELECT OPTION A OR B BELOW*:

- A) PLEASE PROVIDE ASSISTANCE LOCATING AN INFUSION SITE.** PLEASE COORDINATE DIRECTLY WITH: HEALTHCARE PROVIDER PATIENT
 B) ASSISTANCE IS NOT NEEDED. PATIENT WILL BE INFUSED AT: PRESCRIBER'S OFFICE PATIENT'S HOME PREFERRED INFUSION SITE (PLEASE SPECIFY BELOW)

SITE OF CARE NAME	NPI #	TAX ID #
ADDRESS		
CITY	STATE	ZIP
OFFICE CONTACT FOR FOLLOW-UP		PHONE NUMBER

Please see Indications & Important Safety Information on page 4 and full [Prescribing Information and Medication Guide](#) for ULTOMIRIS, including **Boxed WARNING** regarding serious and life-threatening meningococcal infections/sepsis, also available on www.ULTOMIRIS.com.

Please see Indications & Important Safety Information on page 5 and full [Prescribing Information and Medication Guide](#) for SOLIRIS, including **Boxed WARNING** regarding serious and life-threatening meningococcal infections, also available on www.SOLIRIS.net.

PRESCRIBER START FORM - GENERALIZED MYASTHENIA GRAVIS (gMG)



FAX: 1.800.420.5150



MAIL: 100 College Street
New Haven, CT 06510



PHONE: 1.888.765.4747
8:30 AM to 8 PM ET Monday-Friday



EMAIL: OneSource@Alexion.com



Fields in red with asterisks are required.*

PATIENT INFORMATION

PATIENT NAME (FIRST, LAST)*

DATE OF BIRTH (MM/DD/YYYY)*

STEP 6: CLINICAL INFORMATION

CHECK ALL PREVIOUS GENERALIZED MYASTHENIA GRAVIS (gMG) THERAPIES:

- AZATHIOPRINE MYCOPHENOLATE MOFETIL PREDNISONE RITUXIMAB OTHER
 IVIg PLASMAPHERESIS PYRIDOSTIGMINE EFGARTIGIMOD

MGFA CLASSIFICATION: _____

CURRENT MG-ADL SCORE: _____

Abbreviations: IVIg, intravenous immunoglobulin; MG-ADL, Myasthenia Gravis Activities of Daily Living; MGFA, Myasthenia Gravis Foundation of America.

STEP 7: PRESCRIPTION

YOU MAY USE THIS SECTION TO PROVIDE A PRESCRIPTION FOR ULTOMIRIS OR SOLIRIS, OR YOU MAY PROVIDE A SEPARATE PRESCRIPTION.

Rx **ULTOMIRIS** 100 mg/mL HCPCS CODE: J1303 PER UNIT
PATIENT WEIGHT: _____

Rx **SOLIRIS** 10 mg/mL HCPCS CODE: J1300 PER UNIT

LOADING DOSE:

SIG: INFUSE INTRAVENOUSLY _____ mg
ON DAY 0. COVERS THE PATIENT FOR THE
FIRST 2 WEEKS.

OTHER: _____

QTY OF 300 mg/3 mL

VIALS: _____ REFILLS: 0

MAINTENANCE DOSE:

SIG: INFUSE INTRAVENOUSLY _____ mg
EVERY 8 WEEKS. START 2 WEEKS AFTER
COMPLETION OF LOADING DOSE.

OTHER: _____

QTY OF 300 mg/3 mL

VIALS: _____ REFILLS: _____

QTY OF 1100 mg/11 mL

VIALS: _____ REFILLS: _____

LOADING DOSE:

SIG: INFUSE INTRAVENOUSLY _____ mg
WEEKLY FOR THE FIRST 4 WEEKS, FOLLOWED
BY _____ mg FOR THE 5TH WEEK.

OTHER: _____

QTY OF 300 mg/30 mL

VIALS: _____ REFILLS: 0

MAINTENANCE DOSE:

SIG: INFUSE INTRAVENOUSLY _____ mg
EVERY 2 WEEKS. START 2 WEEKS AFTER
THE 5TH WEEK'S DOSE IS COMPLETE.

OTHER: _____

QTY OF 300 mg/30 mL

VIALS: _____ REFILLS: _____



HAS YOUR PATIENT RECEIVED ANY DOSES OF A MENINGOCOCCAL VACCINE OR ANTIBIOTIC PROPHYLAXIS?
IF SO, PLEASE PROVIDE RELEVANT INFORMATION.
See ACIP guidelines below.*

Alexion complement-inhibitor therapies are available only through a restrictive program under a Risk Evaluation and Mitigation Strategy (REMS).
Vaccination dates provided as part of this form are used to confirm vaccination prior to starting treatment.

YES	Antibiotic prophylaxis administered? <input type="checkbox"/> Yes <input type="checkbox"/> No If yes, start date: ____ / ____ / ____		NO
	Patient has received or is scheduled to receive the required vaccinations per ACIP guidelines. Please complete the following information:		
	MenACWY	MenB	
	1st Dose Date: ____ / ____ / ____ <input type="checkbox"/> Menveo <input type="checkbox"/> Menactra <input type="checkbox"/> MenQuadfi	1st Dose Date: ____ / ____ / ____ <input type="checkbox"/> Bexsero <input type="checkbox"/> Trumenba	Patient needs <input type="checkbox"/> VACCINATION SUPPORT from OneSource <input checked="" type="checkbox"/> Sign prescriber certification below <input checked="" type="checkbox"/> Continue to PAGE 3 to fill out a vaccination prescription *
	2nd Dose Date: ____ / ____ / ____ <input type="checkbox"/> Menveo <input type="checkbox"/> Menactra <input type="checkbox"/> MenQuadfi	2nd Dose Date: ____ / ____ / ____ <input type="checkbox"/> Bexsero <input type="checkbox"/> Trumenba	
	<input checked="" type="checkbox"/> Sign prescriber certification below	3rd Dose Date: ____ / ____ / ____ (3rd dose - Trumenba ONLY)	

*The current ACIP guidelines recommend a regimen of MenACWY AND MenB doses prior to starting a complement inhibitor treatment.

*You may also provide a separate prescription.

STEP 8: PRESCRIBER CERTIFICATION

By signing below, I attest that: (i) I am prescribing ULTOMIRIS or SOLIRIS for the patient identified above based on my clinical judgment that it is medically necessary for the diagnosis identified on this form and I will be supervising the patient's treatment; (ii) I am authorized under applicable law to prescribe ULTOMIRIS or SOLIRIS and I have verified and complied with all applicable prescription requirements; (iii) I am authorizing Alexion to forward the patient's prescription to a pharmacy by any means under applicable law; (iv) I am under no obligation to prescribe ULTOMIRIS or SOLIRIS and I have not received, nor will I receive, any benefit from Alexion for prescribing ULTOMIRIS or SOLIRIS; and (v) the information provided on this form is complete, current, and accurate to the best of my knowledge. I also acknowledge that Alexion will use and share the personal data collected about me (as the prescriber) in accordance with the Privacy Notice on the Alexion website at <https://alexion.com/Legal#privacy>.

SIGN ONE*



PRESCRIBER'S SIGNATURE (NO STAMPS) - **DISPENSE AS WRITTEN**

DATE (MM/DD/YYYY)

PRESCRIBER'S SIGNATURE (NO STAMPS) - **MAY SUBSTITUTE**

DATE (MM/DD/YYYY)

Please verify your local prescribing requirements (eg, New York prescribers must provide a separate prescription).

Please see **Indications & Important Safety Information on page 4** and full **Prescribing Information and Medication Guide for ULTOMIRIS**, including **Boxed WARNING** regarding serious and life-threatening meningococcal infections/sepsis, also available on www.ULTOMIRIS.com.

Please see **Indications & Important Safety Information on page 5** and full **Prescribing Information and Medication Guide for SOLIRIS**, including **Boxed WARNING** regarding serious and life-threatening meningococcal infections, also available on www.SOLIRIS.net.

VACCINATION ORDER FORM



FAX: 1.800.420.5150



MAIL: 100 College Street
New Haven, CT 06510



PHONE: 1.888.765.4747
8:30 AM to 8 PM ET Monday-Friday



EMAIL: OneSource@Alexion.com



Personalized Patient Support from Alexion

PATIENT INFORMATION

PATIENT NAME (FIRST, MIDDLE INITIAL, LAST)		PATIENT DATE OF BIRTH (MM/DD/YYYY)	
ADDRESS	CITY	STATE	ZIP
PHONE NUMBER	HEIGHT	WEIGHT	

HEALTHCARE PRESCRIBER INFORMATION

FIRST NAME	LAST NAME	PHONE NUMBER	FAX NUMBER
ADDRESS	CITY	STATE	ZIP
OFFICE CONTACT NAME	NPI		

CLINICAL INFORMATION

Primary Diagnosis Description: Encounter for Immunization **ICD-10 CODE:** Z23

MENINGOCOCCAL VACCINATIONS ARE INDICATED FOR PATIENTS, INCLUDING PEOPLE OVER 25 YEARS OF AGE, WHEN ON A COMPLEMENT INHIBITOR TREATMENT.

The Advisory Committee on Immunization Practices (ACIP) recommends a regimen of MenACWY AND MenB doses prior to starting a complement inhibitor treatment. Vaccines should be initiated at least 2 weeks prior to first dose of Alexion Complement Inhibitor. There are two (2) types of meningococcal vaccines available in the United States.

MenACWY

ONE (1) REQUIRED FROM EACH GROUP

MenB

MenB VACCINES ARE NOT INTERCHANGEABLE. PATIENT MUST RECEIVE THE SAME PRODUCT FOR ALL DOSES DURING A VACCINATION SERIES. FOR THE FULL VACCINE SCHEDULE, INCLUDING THE VACCINATION SCHEDULE FOR CHILDREN ≤10 YEARS OLD, PLEASE REFER TO THE ACIP VACCINE RECOMMENDATIONS OR TO ALEXION MEDICAL INFORMATION.

Two quadrivalent meningococcal conjugate (MenACWY) vaccines are currently licensed and available in the United States.

INDICATE VACCINE THE PATIENT NEEDS TO RECEIVE:

- MenQuadfi** (meningococcal groups A, C, W, and Y polysaccharide tetanus toxoid conjugate vaccine [MenACWY-TT]) 90619
- Menveo** (meningococcal groups A, C, W, and Y oligosaccharide diphtheria CRM conjugate vaccine [MenACWY-CRM]) 907340

Two serogroup B meningococcal (MenB) vaccines are currently licensed and available in the United States.

INDICATE VACCINE THE PATIENT NEEDS TO RECEIVE:

- Bexsero** (MenB-4C) 90620
- Trumenba** (MenB-FHbp) 90621

MenACWY

DOSING SCHEDULE

MenB

- Dose 1:** Day 0
- Dose 2:** At least 8 weeks after Day 0
- Booster dose**

- Dose 1:** Day 0
- Dose 2:** For Bexsero: At least (or greater than or equal to) 1 month after Day 0
For Trumenba: 1-2 months after Day 0
- Dose 3 (Trumenba only):** 6 months after Day 0
- Booster dose**

Per CDC recommendations, those who remain at increased risk need regular booster doses. MenACWY: For children under the age of 7 years, administer a booster dose 3 years after completion of the primary series and every 5 years thereafter. For children 7 years old or older and adults, administer a booster dose 5 years after completion of the primary series and every 5 years thereafter. MenB: Administer a booster dose of vaccine 1 year after series completion and then every 2 to 3 years thereafter.

NOTE: ALL VACCINES LISTED ABOVE ARE ADMINISTERED INTRAMUSCULARLY AT A DOSE OF 0.5 mL

ANCILLARY ORDERS (HOME ADMINISTRATION ONLY - USE AS NEEDED)

Anaphylaxis Kit - The following items will be dispensed:

- Diphenhydramine 50 mg/mL 1 mL vial x 1. Inject 25 mg IM PRN for allergic reaction. May repeat x 1 dose in 15 min PRN if no improvement
- NS 500 mL bag x 1. Infuse 500 mL IV at KVO rate PRN anaphylaxis
- Epinephrine ampule/vial 1 mg/mL (1:1000) 1 mL x 2 ampules/vials. Inject 0.3 mg SQ PRN for adverse reaction. May repeat x 1 dose in 5 to 15 min PRN

General Anaphylaxis Instructions

Administer emergency medications as ordered. Administer epinephrine as above and repeat dose if necessary. Administer injectable diphenhydramine as above and repeat dose if necessary. Place peripheral IV and administer NS. Initiate CPR if needed. Call EMS (activate the emergency medical system). Monitor vital signs—elevate legs if hypotensive. Notify prescriber and Nursing Director or pharmacist.

PRESCRIBER CERTIFICATION

By signing below, I attest that: (i) based on my clinical judgment, the vaccines identified are medically necessary for the patient and diagnosis identified on this form; (ii) I am authorized under applicable law to prescribe the vaccines identified and I have verified and complied with all applicable prescription requirements; (iii) I am authorizing Alexion to forward the patient's prescription to a pharmacy by any means under applicable law; (iv) I am under no obligation to prescribe the vaccines identified and I have not received, nor will I receive, any benefit from Alexion; and (v) the information provided on this form is complete, current, and accurate to the best of my knowledge.



PRESCRIBER SIGNATURE (NO STAMPS) - **DISPENSE AS WRITTEN**

DATE (MM/DD/YYYY)

PRESCRIBER SIGNATURE (NO STAMPS) - **MAY SUBSTITUTE**

DATE (MM/DD/YYYY)

Please verify your local prescribing requirements (eg, New York prescribers must provide a separate prescription)

ALEXION, the Alexion logo, and the OneSource logo are registered trademarks of Alexion Pharmaceuticals, Inc., and OneSource is a trademark of Alexion Pharmaceuticals, Inc.

© 2022, Alexion Pharmaceuticals, Inc. All rights reserved. US/ULT-g/0099 11/22



FAX: 1.800.420.5150



MAIL: 100 College Street
New Haven, CT 06510



PHONE: 1.888.765.4747
8:30 AM to 8 PM ET Monday-Friday



EMAIL: OneSource@Alexion.com

INDICATION & IMPORTANT SAFETY INFORMATION FOR ULTOMIRIS® (ravulizumab-cwvz)

INDICATION

What is ULTOMIRIS?

ULTOMIRIS is a prescription medicine used to treat adults with a disease called generalized Myasthenia Gravis (gMG) who are anti-acetylcholine receptor (AChR) antibody positive.

It is not known if ULTOMIRIS is safe and effective for the treatment of gMG in children.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about ULTOMIRIS?

ULTOMIRIS is a medicine that affects your immune system and can lower the ability of your immune system to fight infections.

- ULTOMIRIS increases your chance of getting serious and life-threatening meningococcal infections that may quickly become life-threatening and cause death if not recognized and treated early.

- You must receive meningococcal vaccines at least 2 weeks before your first dose of ULTOMIRIS if you are not vaccinated.
- If your healthcare provider decided that urgent treatment with ULTOMIRIS is needed, you should receive meningococcal vaccination as soon as possible.
- If you have not been vaccinated and ULTOMIRIS therapy must be initiated immediately, you should also receive 2 weeks of antibiotics with your vaccinations.
- If you had a meningococcal vaccine in the past, you might need additional vaccination. Your healthcare provider will decide if you need additional vaccination.
- Meningococcal vaccines reduce but do not prevent all meningococcal infections. Call your healthcare provider or get emergency medical care right away if you get any of these signs and symptoms of a meningococcal infection: headache with nausea or vomiting, headache and fever, headache with a stiff neck or stiff back, fever, fever and a rash, confusion, muscle aches with flu-like symptoms and eyes sensitive to light.

Your healthcare provider will give you a Patient Safety Card about the risk of meningococcal infection. Carry it with you at all times during treatment and for 8 months after your last ULTOMIRIS dose. It is important to show this card to any healthcare provider or nurse to help them diagnose and treat you quickly.

ULTOMIRIS is only available through a program called the ULTOMIRIS REMS. Before you can receive ULTOMIRIS, your healthcare provider must: enroll in the ULTOMIRIS REMS program; counsel you about the risk of meningococcal infection; give you information and a **Patient Safety Card** about the symptoms and your risk of meningococcal infection (as discussed above); and make sure that you are vaccinated with a meningococcal vaccine, and if needed, get revaccinated with the meningococcal vaccine. Ask your healthcare provider if you are not sure if you need to be revaccinated.

ULTOMIRIS may also increase the risk of other types of serious infections. Call your healthcare provider right away if you have any new signs or symptoms of infection.

Who should not receive ULTOMIRIS?

Do not receive ULTOMIRIS if you have a meningococcal infection or have not been vaccinated against meningococcal infection unless your healthcare provider decides that urgent treatment with ULTOMIRIS is needed.

Before you receive ULTOMIRIS, tell your healthcare provider about all of your medical conditions, including if you: have an infection or fever, are pregnant or plan to become pregnant, and are breastfeeding or plan to breastfeed. It is not known if ULTOMIRIS will harm your unborn baby or if it passes into your breast milk. You should not breastfeed during treatment and for 8 months after your final dose of ULTOMIRIS.

Tell your healthcare provider about all the vaccines you receive and medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements which could affect your treatment.

What are the possible side effects of ULTOMIRIS?

ULTOMIRIS can cause serious side effects including infusion-related reactions. Symptoms of an infusion-related reaction with ULTOMIRIS may include lower back pain, tiredness, feeling faint, discomfort in your arms or legs, bad taste, or drowsiness. Stop treatment of ULTOMIRIS and tell your healthcare provider or nurse right away if you develop these symptoms, or any other symptoms during your ULTOMIRIS infusion that may mean you are having a serious infusion reaction, including: chest pain, trouble breathing or shortness of breath, swelling of your face, tongue, or throat, and feel faint or pass out.

The most common side effects of ULTOMIRIS in people with gMG are diarrhea and upper respiratory tract infections.

Tell your healthcare provider about any side effect that bothers you or that does not go away. These are not all the possible side effects of ULTOMIRIS. For more information, ask your healthcare provider or pharmacist. Call your healthcare provider right away if you miss an ULTOMIRIS infusion or for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Please see full Prescribing Information and Medication Guide for ULTOMIRIS, including **Boxed WARNING** regarding serious and life-threatening meningococcal infections/sepsis, also available on www.ULTOMIRIS.com.



FAX: 1.800.420.5150



MAIL: 100 College Street
New Haven, CT 06510



PHONE: 1.888.765.4747
8:30 AM to 8 PM ET Monday-Friday



EMAIL: OneSource@Alexion.com

INDICATION & IMPORTANT SAFETY INFORMATION FOR SOLIRIS® (eculizumab)

INDICATION

What is SOLIRIS?

SOLIRIS is a prescription medicine used to treat adults with a disease called generalized myasthenia gravis (gMG) who are anti-acetylcholine receptor (AChR) antibody positive.

It is not known if SOLIRIS is safe and effective in children with gMG.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about SOLIRIS?

SOLIRIS is a medicine that affects your immune system and can lower the ability of your immune system to fight infections.

- **SOLIRIS increases your chance of getting serious and life-threatening meningococcal infections that may quickly become life-threatening and cause death if not recognized and treated early.**

1. You must receive meningococcal vaccines at least 2 weeks before your first dose of SOLIRIS if you are not vaccinated.
2. If your doctor decided that urgent treatment with SOLIRIS is needed, you should receive meningococcal vaccination as soon as possible.
3. If you have not been vaccinated and SOLIRIS therapy must be initiated immediately, you should also receive two weeks of antibiotics with your vaccinations.
4. If you had a meningococcal vaccine in the past, you might need additional vaccination. Your doctor will decide if you need additional vaccination.
5. Meningococcal vaccines reduce but do not prevent all meningococcal infections. Call your doctor or get emergency medical care right away if you get any of these signs and symptoms of a meningococcal infection: headache with nausea or vomiting, headache and fever, headache with a stiff neck or stiff back, fever, fever and a rash, confusion, muscle aches with flu-like symptoms, and eyes sensitive to light.

Your doctor will give you a Patient Safety Card about the risk of meningococcal infection. Carry it with you at all times during treatment and for 3 months after your last SOLIRIS dose. It is important to show this card to any doctor or nurse to help them diagnose and treat you quickly.

SOLIRIS is only available through a program called the SOLIRIS REMS. Before you can receive SOLIRIS, your doctor must enroll in the SOLIRIS REMS program; counsel you about the risk of meningococcal infection; give you information and a **Patient Safety Card** about the symptoms and your risk of meningococcal infection (as discussed above); and make sure that you are vaccinated with the meningococcal vaccine and, if needed, get revaccinated with the meningococcal vaccine. Ask your doctor if you are not sure if you need to be revaccinated.

SOLIRIS may also increase the risk of other types of serious infections. Certain people may be at risk of serious infections with gonorrhea. Certain fungal infections (*Aspergillus*) may occur if you take SOLIRIS and have a weak immune system or a low white blood cell count.

Who should not receive SOLIRIS?

Do not receive SOLIRIS if you have a meningococcal infection or have not been vaccinated against meningitis infection unless your doctor decides that urgent treatment with SOLIRIS is needed.

Before you receive SOLIRIS, tell your doctor about all of your medical conditions, including if you: have an infection or fever, are pregnant or plan to become pregnant, and are breastfeeding or plan to breastfeed. It is not known if SOLIRIS will harm your unborn baby or if it passes into your breast milk.

Tell your doctor about all the vaccines you receive and medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements which could affect your treatment. It is important that you have all recommended vaccinations before you start SOLIRIS, receive 2 weeks of antibiotics if you immediately start SOLIRIS, and stay up-to-date with all recommended vaccinations during treatment with SOLIRIS.

What are the possible side effects of SOLIRIS?

SOLIRIS can cause serious side effects including serious infusion-related reactions. Tell your doctor or nurse right away if you get any of these symptoms during your SOLIRIS infusion: chest pain, trouble breathing or shortness of breath, swelling of your face, tongue, or throat, and feel faint or pass out. If you have an infusion-related reaction to SOLIRIS, your doctor may need to infuse SOLIRIS more slowly, or stop SOLIRIS.

The most common side effects in people with gMG treated with SOLIRIS include: muscle and joint (musculoskeletal) pain.

Tell your doctor about any side effect that bothers you or that does not go away. These are not all the possible side effects of SOLIRIS. For more information, ask your doctor or pharmacist. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit MedWatch, or call 1-800-FDA-1088.

Please see full [Prescribing Information](#) and [Medication Guide](#) for SOLIRIS, including **Boxed WARNING** regarding serious and life-threatening meningococcal infections, also available on www.SOLIRIS.net.

This material is intended only for residents of the United States.

PATIENT SERVICES ENROLLMENT FORM



EMAIL: OneSource@Alexion.com



PHONE: 1.888.765.4747 8:30 AM to 8 PM ET Monday-Friday



FAX: 1.800.420.5150



MAIL: 100 College St., New Haven, CT 06510



OneSource™ is a complimentary, personalized patient support program offered by Alexion. It's designed to support patients' specific needs throughout treatment. For more information, visit www.AlexionOneSource.com.



INSTRUCTIONS FOR PATIENTS:

To enroll in OneSource, please follow these steps:

- 1 Complete all the required information (in red) on **this page** and read the Authorization to Share Health Information on **the next page**
- 2 Sign the Authorization to Share Health Information section on **this page**
- 3 Email or fax **this page** and **copies of the front and back of your medical insurance and pharmacy coverage cards** to OneSource (see the email address and fax number above)

Be sure to complete all required fields and sign and date the form. If information is incomplete, it could delay our ability to enroll you in OneSource. OneSource can start offering you personalized support once you submit this form fully and correctly completed.

Fields in red with asterisks are required.* **Contact OneSource if you have any questions while completing the form.**

PATIENT INFORMATION

PATIENT NAME (FIRST, MIDDLE INITIAL, LAST)*		DATE OF BIRTH (MM/DD/YYYY)*	GENDER: <input type="checkbox"/> MALE <input type="checkbox"/> FEMALE <input type="checkbox"/> NON-BINARY PREFER TO SELF-DESCRIBE:
ADDRESS*			
CITY*		STATE*	ZIP*
PRIMARY PHONE NUMBER* <input type="checkbox"/> MOBILE <input type="checkbox"/> HOME	OK TO SEND A TEXT MESSAGE? <input type="checkbox"/> YES <input type="checkbox"/> NO OK TO LEAVE A PHONE MESSAGE? <input type="checkbox"/> YES <input type="checkbox"/> NO		
PATIENT DIAGNOSIS			

PREFERRED LANGUAGE <input type="checkbox"/> ENGLISH <input type="checkbox"/> SPANISH <input type="checkbox"/> OTHER _____	PATIENT EMAIL <input type="checkbox"/> NONE	
LEGAL PATIENT REPRESENTATIVE* (REQUIRED IF A PATIENT IS A MINOR)	RELATIONSHIP TO PATIENT	EMAIL
NAME: _____ PHONE: _____		
DESIGNATED CARE PARTNER	RELATIONSHIP TO PATIENT	EMAIL
NAME: _____ PHONE: _____		

PRESCRIBING PHYSICIAN'S INFORMATION

PROVIDER NAME	PROVIDER PHONE NUMBER	PROVIDER EMAIL
---------------	-----------------------	----------------

AUTHORIZATION TO SHARE HEALTH INFORMATION

By signing below, I acknowledge that I have read and agree to the Authorization to Share Health Information terms on the next page.

SIGN HERE*



SIGNATURE OF PATIENT OR LEGALLY AUTHORIZED REPRESENTATIVE

DATE (MM/DD/YYYY)

CONSENT FOR COPAY PROGRAM (OPTIONAL)

By signing below, I acknowledge that I have read and agree to the Alexion OneSource CoPay Program terms and conditions available at <https://alexiononesource.com/CoPay> or on request by contacting OneSource at 1.888.765.4747.

SIGNATURE OF PATIENT OR LEGALLY AUTHORIZED REPRESENTATIVE

DATE (MM/DD/YYYY)

CONSENT FOR AUTOMATED TEXT COMMUNICATIONS (OPTIONAL)

By signing below, I give Alexion and companies working at Alexion's direction permission to use automated text (SMS) messages to provide patient support services and to provide information to me about Alexion products, services, programs, or other topics that Alexion thinks may interest me. I understand that (i) I am not required to consent to receiving text messages as a condition of any purchase of Alexion products or enrollment in these programs; (ii) my telecommunication services provider may charge me for any text messages that I receive from Alexion; and (iii) I may opt out of receiving automated text messages from Alexion at any time without affecting my enrollment in these programs.

SIGNATURE OF PATIENT OR LEGALLY AUTHORIZED REPRESENTATIVE

DATE (MM/DD/YYYY)

PATIENT SERVICES ENROLLMENT FORM



EMAIL: OneSource@Alexion.com



PHONE: 1.888.765.4747 8:30 AM to 8 PM ET Monday-Friday



FAX: 1.800.420.5150



MAIL: 100 College St., New Haven, CT 06510



AUTHORIZATION TO SHARE HEALTH INFORMATION

Alexion Pharmaceuticals, Inc. (“Alexion”) offers patient services including educational resources, case management support, and financial assistance for eligible patients.

By signing the prior page, I give permission for my healthcare providers, health plans, other insurance programs, pharmacies, and other healthcare service providers (“My Healthcare Entities”) to share information, including protected health information relating to my medical condition, treatment, and health insurance coverage (collectively “My Information”) with Alexion and companies working at its direction so that Alexion may use and disclose My Information to:

- review my eligibility for benefits for treatment with an Alexion product;
- coordinate treatment with an Alexion product, as well as related services, such as arranging home infusion services or vaccine services;
- access my credit information and information from other sources to estimate my income, if needed, to assess eligibility for financial assistance programs;
- remove identifiers from My Information and combine such resulting information with other information for research, regulatory submissions, business improvement projects, and publication purposes; and
- contact me about market research or clinical studies, provide me with educational and promotional materials, or otherwise contact me about Alexion products, services, programs, or other topics that Alexion thinks may interest me.

I understand that My Healthcare Entities may receive payment from Alexion in exchange for sharing My Information.

I understand that My Information is also subject to the Alexion Privacy Notice available at <https://alexion.com/Legal#privacy>, and that the Alexion Privacy Notice provides additional information about Alexion’s privacy practices and the rights that may be available to me. Although Alexion has implemented privacy and security controls designed to help protect My Information, I understand that once My Information has been disclosed to Alexion, the Health Insurance Portability and Affordability Act (“HIPAA”) may not apply and may no longer protect the information.

I understand that I may refuse to sign this Authorization and that My Healthcare Entities may not condition treatment, payment, enrollment, or eligibility for benefits on whether I sign this Authorization. I also understand that if I do not sign this Authorization, I will not be able to receive support through the Alexion OneSource™ Patient Support Program.

This Authorization expires ten (10) years from the date next to my signature, unless I cancel/revoke it sooner, or unless a shorter time frame is required by applicable law.

I understand that I may revoke my authorization, or unsubscribe or modify the services I receive, at any time by mailing a letter to Alexion OneSource Patient Support Program, 100 College Street, New Haven, CT 06510 or by emailing OneSource@Alexion.com. I also understand that modifying my authorization will not affect any use or disclosure of My Information that occurred before Alexion received notice of my cancellation. I also understand I have a right to receive a copy of this Authorization after it is signed and can request a copy at any time by contacting OneSource at 1.888.765.4747.

OneSource Services

Alexion services and support are subject to change. Participation is voluntary, and person(s) may be removed from Alexion services for code of conduct violations.

This material is intended only for residents of the United States.

ALEXION, the Alexion logo, and the OneSource logo are registered trademarks of Alexion Pharmaceuticals, Inc., and OneSource is a trademark of Alexion Pharmaceuticals, Inc.

© 2022, Alexion Pharmaceuticals, Inc. All rights reserved. US/ULT-g/0099 11/22

